

•
•
•
•
•
•
•

DISCLAIMER

The information contained in this document is restricted and it is intended for professional use by licensed health care providers. The statements made should not be construed as a claim, and does not represent any particular product procedure or advice as constituting a specific cure, whether palliative or ameliorative. Procedures and products mentioned are designed to support the client's health and may be considered as support for other procedures deemed necessary by the practitioner.

Although the supplements, herbs and homeopathics referenced in this document are made by specific manufacturers, the opinions expressed are those of the author and are not those of any of the manufacturers or their employees. This document is not designed to replace medical advice or medical treatments, but to be used adjunctively in the care and support of health and vitality.



SUPPLEMENTAL PROTOCOLS

RENAL AND BLADDER CONDITIONS

Kidney Support

Primary Support

(MULTIPLE NUTRIENTS FOR SUPPORTING RENAL FUNCTION-NON-GLANDULAR)

Recommendation: 2-4 tablets one to four times daily

Secondary Support

(CULTURE OF BEET JUICE CONTAINING ARGINASE)

Recommendation: 1-4 tablets one to four times daily

(KIDNEY TISSUE)

Recommendation: 1-2 tablets one to four times daily

Kidney Stones

Primary Support for 7-day Stone Flush

(NUTRITIONAL PHOSPHORUS)

Recommendation: 1 bottle in 4 oz. tomato juice once daily

(MULTIPLE NUTRIENTS FOR SUPPORTING RENAL FUNCTION-NON-GLANDULAR)

Recommendation: 2-4 tablets two to four times daily

Secondary Support

(PYRIDOXAL-5-PHOSPHATE)

Recommendation: 1-2 tablets two to four times daily

•
•
•
•
•
•
•

(MAGNESIUM)

Recommendation: 4-8 tablets at bed time to bowel tolerance

Continuing support after the flush:

(MULTIPLE NUTRIENTS FOR SUPPORTING RENAL FUNCTION-NON-GLANDULAR)

Recommendation: 1-2 tablets two to four times daily

If average urinary pH < 5.5 add:

(POTASSIUM)

Recommendation: 1-2 tablets two to four times daily

If average urinary pH > 6.5 add:

(NUTRITIONAL PHOSPHORUS)

Recommendation: 1 dropper once or twice daily

If stones are carbonate:

(NUTRITIONAL PHOSPHORUS)

Recommendation: 1 dropper once or twice daily

If stones are oxalate:

(NUTRITIONAL PHOSPHORUS)

Recommendation: 1 dropper once or twice daily



If stones are urate:

(FOLIC ACID)
Recommendation: 1-2 tablets two to four times daily

(POTASSIUM)
Recommendation: 1-2 tablets two to four times daily

Secondary Support

(PYRIDOXAL-5-PHOSPHATE)
Recommendation: 1-2 tablets two to four times daily

(MAGNESIUM)
Recommendation: 4-8 tablets at bed time to bowel tolerance

Bladder Infections

Primary Support

(MULTIPLE NUTRIENTS FOR SUPPORTING RENAL FUNCTION-NON-GLANDULAR)
Recommendation: 2-4 tablets two to four times daily

(MICRO EMULSIFIED OREGANO OIL)
Recommendation: 2-4 tablets two to four times daily

Secondary Support

(LARCH ARABINOGALACTANS)
Recommendation: 1 teaspoon once or twice daily

(NUTRIENTS THAT SUPPORT AGAINST BACTERIA)
Recommendation: 2-4 tablets two to four times daily

•
•
•
•
•
•
•
•

Leaky Bladder / Interstitial cystitis

Primary Support

(MULTIPLE NUTRIENTS FOR SUPPORTING RENAL FUNCTION-NON-GLANDULAR)

Recommendation: 2-4 tablets two to four times daily

(NATURALLY OCCURRING THIAMINE)

Recommendation: 2-4 tablets two to four times daily

Secondary Support

(EMULSIFIED VITAMIN A DROPS)

Recommendation: 2-4 tablets two to four times daily

(CULTURE OF BEET JUICE CONTAINING ARGINASE)

Recommendation: 2-4 tablets two to four times daily

If average urinary pH < 5.5:

(POTASSIUM)

Recommendation: 1-2 tablets two to four times daily

If average urinary pH > 6.5:

(NUTRITIONAL PHOSPHORUS)

Recommendation: 1 dropper once or twice daily